

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
11 May 2006 (11.05.2006)

PCT

(10) International Publication Number  
**WO 2006/050122 A1**

(51) International Patent Classification:  
C08G 69/10 (2006.01)

(74) Agent: THALLEMER, John, D.; Novartis, Corporate Intellectual Property Department, One Health Plaza, Bldg. 104, East Hanover, New Jersey 07936 (US).

(21) International Application Number:  
PCT/US2005/039003

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date: 27 October 2005 (27.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/623,346 29 October 2004 (29.10.2004) US  
60/651,372 9 February 2005 (09.02.2005) US  
60/651,535 9 February 2005 (09.02.2005) US  
60/723,901 5 October 2005 (05.10.2005) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): SANDOZ AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(71) Applicant (for AT only): NOVARTIS PHARMA GmbH [AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): RAY, Anup, Kumar [IN/US]; 215 Aspen Knolls Way, Staten Island, New York 10312 (US). PATEL, Hiren, Kumar, V. [US/US]; 9 Wadsworth Lane, Fords, New Jersey 08863 (US). LUDSCHER, Johannes [AT/AT]; Kleinsöhl 101, A-6252 Breitenbach (AT). ANBAZHAGAN, Mariappan [IN/US]; 307 Northumberland Way, Monmouth Junction, New Jersey 08852 (US). PATEL, Mahendra, R. [US/US]; 5 Belair Court, Milltown, New Jersey 08850 (US). MACHER, Ingolf [AT/AT]; Moosweg 19, A-6300 Woergl (AT).

**Declaration under Rule 4.17:**

— as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))

**Published:**

— with international search report  
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PROCESSES FOR PREPARING GLATIRAMER

(57) Abstract: The present invention relates to processes for preparing a polypeptide or pharmaceutically acceptable salt thereof comprising L-tyrosine, L-alanine, L-glutamate and L-lysine. The polypeptide or pharmaceutically acceptable salt thereof is preferably glatiramer acetate.

BEST AVAILABLE COPY

WO 2006/050122 A1

## PROCESSES FOR PREPARING GLATIRAMER

Field of the Invention

The present invention provides processes for preparing a polypeptide or pharmaceutically acceptable salt thereof. More specifically, the invention provides processes for preparing glatiramer acetate.

Background of the Invention

COPAXONE® is the trade name of glatiramer acetate, an FDA approved drug for the treatment of multiple sclerosis. COPAXONE® is also known as Copolymer-1. The COPAXONE® label discloses that COPAXONE® consists of the acetate salts of synthetic polypeptides, containing four naturally-occurring amino acids: L-glutamic acid, L-alanine, L-tyrosine and L-lysine with an average molar fraction of 0.141, 0.427, 0.095 and 0.338, respectively, and has a weight average molecular weight of 4.7-11.0 kilodaltons (kDa). COPAXONE® comprises a mixture of polypeptides having different molecular weights and sequences. The structural formula of COPAXONE® is:



COPAXONE® is a white to off-white, sterile, lyophilized powder containing 20 mg glatiramer acetate and 40 mg of mannitol. It is supplied in single use vials for subcutaneous administration after reconstitution with sterile water.

Processes for preparing Copolymer-1 or glatiramer acetate have been described in U.S. Patent Nos. 3,849,550; 5,800,808; 5,981,589; 6,048,898; 6,054,430; 6,342,476; and 6,362,161. The process for the synthesis of glatiramer acetate is based on the polymerization of *N*-carboxyanhydrides of tyrosine, alanine,  $\gamma$ -benzyl glutamate and *N*<sup>ε</sup>-trifluoroacetyl lysine in anhydrous dioxane at room temperature using diethylamine as initiator, to form a protected polypeptide. The deblocking of the  $\gamma$ -benzyl groups (first deprotection) is accomplished by stirring the protected polypeptide in hydrogen bromide/acetic acid at room temperature. These conditions also facilitate the cleavage of the copolymer. The next step is the removal of the *N*<sup>ε</sup>-trifluoroacetyl groups (second deprotection) of the copolymer by treatment with 1 M piperidine. In the final steps, glatiramer acetate is obtained by purification of the copolymer through dialysis, followed by treatment

## PROCESSES FOR PREPARING GLATIRAMER

Field of the Invention

The present invention provides processes for preparing a polypeptide or pharmaceutically acceptable salt thereof. More specifically, the invention provides processes for preparing glatiramer acetate.

Background of the Invention

COPAXONE® is the trade name of glatiramer acetate, an FDA approved drug for the treatment of multiple sclerosis. COPAXONE® is also known as Copolymer-1. The COPAXONE® label discloses that COPAXONE® consists of the acetate salts of synthetic polypeptides, containing four naturally-occurring amino acids: L-glutamic acid, L-alanine, L-tyrosine and L-lysine with an average molar fraction of 0.141, 0.427, 0.095 and 0.338, respectively, and has a weight average molecular weight of 4.7-11.0 kilodaltons (kDa). COPAXONE® comprises a mixture of polypeptides having different molecular weights and sequences. The structural formula of COPAXONE® is:



COPAXONE® is a white to off-white, sterile, lyophilized powder containing 20 mg glatiramer acetate and 40 mg of mannitol. It is supplied in single use vials for subcutaneous administration after reconstitution with sterile water.

Processes for preparing Copolymer-1 or glatiramer acetate have been described in U.S. Patent Nos. 3,849,550; 5,800,808; 5,981,589; 6,048,898; 6,054,430; 6,342,476; and 6,362,161. The process for the synthesis of glatiramer acetate is based on the polymerization of *N*-carboxyanhydrides of tyrosine, alanine,  $\gamma$ -benzyl glutamate and *N*<sup>ε</sup>-trifluoroacetyl lysine in anhydrous dioxane at room temperature using diethylamine as initiator, to form a protected polypeptide. The deblocking of the  $\gamma$ -benzyl groups (first deprotection) is accomplished by stirring the protected polypeptide in hydrogen bromide/acetic acid at room temperature. These conditions also facilitate the cleavage of the copolymer. The next step is the removal of the *N*<sup>ε</sup>-trifluoroacetyl groups (second deprotection) of the copolymer by treatment with 1 M piperidine. In the final steps, glatiramer acetate is obtained by purification of the copolymer through dialysis, followed by treatment

with acetic acid to form the acetate salt and by another purification by dialysis against water. Thus, these prior art processes involve the polymerization of four *N*-carboxyanhydrides, two deprotection steps, two purification steps and one acetate salt formation step.

U.S. Patent No. 6,620,847 describes a process for preparing Copolymer-1 which involves treating trifluoroacetyl Copolymer-1 with aqueous piperidine to form a solution of Copolymer-1 and purifying Copolymer-1.

U.S. Patent Application Publication No. 2004/0091956 describes a three-step process for preparing glatiramer acetate. The process involves polymerization of a mixture of the *N*-carboxyanhydrides of L-alanine, L-tyrosine, protected L-glutamate and protected L-lysine, to obtain a protected polypeptide or salt thereof; and deprotection of the protected polypeptide or salt thereof by either palladium catalytic transfer hydrogenation or palladium catalytic hydrogenation under hydrogen pressure.

#### Summary of the Invention

The invention provides a process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:

- (i) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine; *N*-carboxyanhydride of protected L-glutamate; and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof; and
- (ii) adding an acid to the protected polypeptide formed in Step (i) to form a polypeptide or a pharmaceutically acceptable salt thereof, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety.

The invention provides a process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises treating a protected polypeptide with an aqueous solution of an alkali or alkaline earth metal hydroxide to form a polypeptide or a pharmaceutically acceptable salt thereof.

The invention provides a process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:

- (a)<sup>2</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
- (b)<sup>2</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>2</sup> and a solvent, to form a product; and
- (c)<sup>2</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, with the product formed in Step (b)<sup>2</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

The invention provides a process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:

- (a)<sup>3</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
- (b)<sup>3</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>3</sup> and a solvent, to form a product; and
- (c)<sup>3</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, with the product formed in Step (b)<sup>3</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

#### Description of the Invention

The present invention relates to processes for preparing a polypeptide or pharmaceutically acceptable salt thereof comprising L-tyrosine, L-alanine, L-glutamate and L-lysine. The polypeptide or pharmaceutically acceptable salt thereof is preferably glatiramer acetate.

## I. Acid Hydrolysis Process.

In one embodiment of the invention, the process comprises:

- (i) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof; and
- (ii) adding an acid to the protected polypeptide formed in Step (i) to form a polypeptide or a pharmaceutically acceptable salt thereof, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety.

In one embodiment of the invention, the process comprises:

- (a) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate, and mixtures thereof;
- (b) adding an acid to the protected glatiramer formed in Step (a) to form a glatiramer, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety; and
- (c) treating the glatiramer formed in Step (b) with acetic acid to form glatiramer acetate.

In the polymerizing step of the processes of the invention, Step (i), Step (a), Step (a)<sup>1</sup>, Step (a)<sup>1'</sup>, Step (a)<sup>2</sup>, Step (a)<sup>2'</sup>, Step (a)<sup>3</sup> and Step (a)<sup>3'</sup>, the mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, are preferably polymerized at a temperature of from about 10°C to about 40°C, more preferably about 20°C to about 30°C. The polymerization reaction preferably takes place for a period of from about 2 hours to about 80 hours, more preferably from about 20 hours to about 50 hours. Most preferably, the polymerization reaction takes place for a period of about 24 hours at a temperature of about 25°C.

The polar aprotic solvent is preferably selected from tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane. Most preferably, the polar aprotic solvent is 1,4-dioxane. A mixture of polar aprotic solvents may also be used.

The initiator used in Step (i), Step (a), Step (a)<sup>1</sup>, Step (a)<sup>1'</sup>, Step (a)<sup>2</sup>, Step (a)<sup>2'</sup>, Step (a)<sup>3</sup> and Step (a)<sup>3'</sup>, of the processes of the invention may be any alkylamine initiator, such as a dialkyl or a trialkylamine. Each of the alkyl groups preferably has 1-6 carbon atoms. A preferred alkylamine initiator is diethylamine. Preferably, the diethylamine is present in an amount of from about 0.001 weight percent (wt. %) to about 2 wt. %, more preferably from about 0.01 wt. % to about 0.02 wt. %, based on the weight of the mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine.

In one embodiment of the invention, water is added to the polymerization mixture following polymerization. The addition of water results in precipitation of the protected polypeptide. The water is preferably removed from the mixture containing water and protected polypeptide by vacuum filtration and the recovered protected polypeptide is dried. Methods of drying are known to those skilled in the art, such as vacuum drying.

In the deprotecting step, Step (ii), of the process of the invention, an acid is added to the protected polypeptide which is formed in Step (i) to form a polypeptide or a pharmaceutically acceptable salt thereof. The acid cleaves the  $\gamma$ -*p*-methoxybenzyl group or  $\gamma$ -benzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety. In addition, the acid cleaves the amide bonds of the polypeptide or pharmaceutically acceptable salt thereof forming heterogenous polypeptide fragments.

In the deprotecting step, Step (b), of the process of the invention, an acid is added to the protected glatiramer formed in Step (a) to form a glatiramer. The acid cleaves the  $\gamma$ -*p*-methoxybenzyl group or the  $\gamma$ -benzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety. In addition, the acid cleaves the amide bonds of the glatiramer forming heterogenous glatiramer fragments.

Suitable acids include, but are not limited to, acetic acid, hydrochloric acid, hydrogen bromide, hydrogen fluoride, methane sulfonic acid, trifluoromethane sulfonic acid, phosphoric acid, trifluoroacetic acid and sulfuric acid. A mixture of acids may also be used. Preferred acids are selected from trifluoroacetic acid, a mixture of acetic acid and hydrochloric acid, a

mixture of acetic acid and hydrogen bromide and a mixture of acetic acid and sulfuric acid. The acid may be added in the form of an aqueous solution.

The acid is preferably present in an amount of from about 0.1 wt. % to about 100 wt. %, more preferably from about 1 wt. % to about 10 wt. %, based on the weight of the protected polypeptide or protected glatiramer. Most preferably, the acid is present in an amount of from about 2 wt. % to about 6 wt. %, based on the weight of the protected polypeptide or protected glatiramer.

The temperature of the reaction medium during addition of the acid is preferably from about 10°C to about 40°C, more preferably 15°C to about 30°C. The acid is preferably added over a period of time from about 1 hour to about 30 hours, with stirring. Most preferably, the acid is added to the protected polypeptide or protected glatiramer at a temperature of about 25°C for a period of from about 1 hour to about 8 hours, with stirring.

Excess acid is preferably removed from the reaction mixture by purging the reaction mixture with nitrogen, lyophilization, or by means of a rotary evaporator under vacuum to obtain a deprotected polypeptide in solid form. However, other separation techniques known to those skilled in the art may also be used.

The deprotected polypeptide or deprotected glatiramer in the form of a free base or acid addition salt is preferably dissolved in water or an aqueous acetic acid solution. Undesired low molecular weight polypeptide or glatiramer fragments, i.e., less than about 2 kDa, and high molecular weight polypeptide or glatiramer fragments, i.e., greater than about 40 kDa, are preferably removed by such methods as dialysis or diafiltration. Preferred membranes include Visking partially permeable cellulose membranes, such as a Size 6 membrane having a molecular weight cut-off of 12-14 kDa, available from Medicell International Ltd., and tangential flow filtration (TFF) membranes, such as a Pellicon XL PLCCC 10 (50 cm<sup>2</sup>) or PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. In a preferred embodiment of the invention, the deprotected polypeptide or deprotected glatiramer is dissolved in water and subjected to dialysis, followed by dialysis in aqueous acetic acid solution.

The present inventors have determined that the desired molecular weight polypeptide or a pharmaceutically acceptable salt thereof may be controlled by dilution, concentration of the acid added in Step (ii) or Step (b), and/or time.



In one embodiment of the invention, water is removed from the deprotected polypeptide. A preferred method of removal is lyophilization. In lyophilization, the solution is frozen and placed under vacuum so that the water (ice) vaporizes in the vacuum (sublimes) without melting and the non-water components (deprotected polypeptide and residual salt) are left behind in an undamaged state, i.e., without chemical decomposition. The dried product of lyophilization contains the deprotected polypeptide and residual salt.

In one embodiment of the invention, the deprotected polypeptide is treated with glacial acetic acid to form glatiramer acetate salt. The glatiramer acetate salt is collected preferably by lyophilization to yield a glatiramer acetate salt product.

## II. Phase Transfer Process.

In one embodiment of the invention, the process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, comprises treating a protected polypeptide with an aqueous solution of an alkali or alkaline earth metal hydroxide to form a polypeptide or a pharmaceutically acceptable salt thereof.

In one embodiment of the invention, the process comprises:

- (a)<sup>1</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide; and
- (b)<sup>1</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected polypeptide formed in Step (a)<sup>1</sup> to form a polypeptide or a pharmaceutically acceptable salt thereof.

In one embodiment of the invention, the process comprises:

- (a)<sup>1'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a  $\gamma$ -benzyl L-glutamate L-glutamate and *N*-carboxyanhydride of N<sup>E</sup>-trifluoroacetyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer;
- (b)<sup>1'</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected glatiramer formed in Step (a)<sup>1'</sup> to form a glatiramer; and
- (c)<sup>1'</sup> treating the glatiramer with acetic acid to form glatiramer acetate.

In the deprotecting steps, Step (b)<sup>1</sup> and Step (b)<sup>1'</sup>, of the process of the invention, an aqueous solution of an alkali or alkaline earth metal hydroxide is added to the protected polypeptide formed in Step (a)<sup>1</sup> or to the protected glatiramer formed in Step (a)<sup>1'</sup>. In Step (b)<sup>1</sup> and Step (b)<sup>1'</sup>, the protecting groups on the glutamic acid moiety, i.e., the  $\gamma$ -benzyl group, and the protecting groups on the lysine moiety, i.e., *N*<sup>ε</sup>-trifluoroacetyl group, are removed. In addition, the alkali or alkaline earth metal hydroxide cleaves the amide bonds of the polypeptide or glatiramer forming heterogenous polypeptide or glatiramer fragments.

While not wishing to be bound by any particular theory, the present inventors believe that the removal of the protecting groups from the protected polypeptide or protected glatiramer results in a phase transfer of the deprotected polypeptide or deprotected glatiramer from the organic phase to the aqueous phase.

In the absence of a buffer in Step (b)<sup>1</sup> or Step (b)<sup>1'</sup>, the pH during Step (b)<sup>1</sup> and Step (b)<sup>1'</sup>, is generally about 13 to about 14 after the addition of the aqueous solution of an alkali or alkaline earth metal hydroxide. In the presence of a buffer in Step (b)<sup>1</sup> or Step (b)<sup>1'</sup>, the pH is generally about 8 to about 12. The buffer may be added or formed *in situ*. A preferred buffer is an acetate buffer such as sodium acetate.

The alkali or alkaline earth metal hydroxide is preferably selected from calcium hydroxide, lithium hydroxide, magnesium hydroxide, potassium hydroxide and sodium hydroxide. More preferably, the alkali or alkaline earth metal hydroxide is sodium hydroxide. A combination of alkali or alkaline earth metal hydroxides may also be used.

The alkali or alkaline earth metal hydroxide is preferably present in an amount of from about 0.1 wt. % to about 400 wt. %, more preferably from about 10 wt. % to about 300 wt. %, based on the weight of the protected polypeptide or protected glatiramer. Most preferably, the alkali or alkaline earth metal hydroxide is present in an amount of from about 140 wt. % to about 260 wt. %, based on the weight of the protected polypeptide or protected glatiramer.

An aqueous solution of an alkali or alkaline earth metal hydroxide is added to the protected polypeptide or protected glatiramer preferably at a temperature of from about -78°C to about 40°C, more preferably -25°C to about 30°C, for a period of time preferably from about 1 hour to about 30 hours. Most preferably, an aqueous solution of an alkali or alkaline earth metal hydroxide is added to the protected polypeptide or protected glatiramer at a temperature from about -10°C to about 10°C, e.g., 0°C, for a period of from about 1 hour to about 8 hours, with stirring. The addition of an aqueous solution of an alkali or alkaline

earth metal hydroxide results in a phase separation wherein an organic phase and an aqueous phase are formed.

The organic phase substantially contains the polar aprotic solvent and the protected polypeptide or protected glatiramer. The aqueous phase substantially contains water, the alkali or alkaline earth metal hydroxide, and the deprotected polypeptide or deprotected glatiramer in the form of a free base. The aqueous phase and the organic phase are preferably separated by using a centrifuge and decanting the organic phase.

An additional advantage of the process of the invention is that the addition of the alkali or alkaline earth metal hydroxide causes deprotection of the protected polypeptide to form a deprotected polypeptide, or deprotection of the protected glatiramer to form a deprotected glatiramer. In addition, the alkali or alkaline earth metal hydroxide causes cleavage of the amide bonds in the deprotected polypeptide or deprotected glatiramer to form polypeptide or glatiramer fragments.

The aqueous phase is preferably treated with an organic or mineral acid to achieve a pH of about 7 to about 8. Such organic or mineral acids to adjust the pH are well-known to those skilled in the art and include, but are not limited to, acetic acid, formic acid, oxalic acid and hydrochloric acid. Dilute hydrochloric acid is preferred. The amount of organic or mineral acid used is preferably an equivalent amount based on the amount of the alkali or alkaline earth metal hydroxide added during Step (b)<sup>1</sup> or Step (b)<sup>1'</sup> which is sufficient to produce a pH of about 7 to about 8.

Undesired low molecular weight polypeptide or glatiramer fragments, i.e., less than about 2 kDa, and high molecular weight polypeptide or glatiramer fragments, i.e., greater than about 40 kDa, are preferably removed by such methods as dialysis or diafiltration. Preferred membranes include Visking partially permeable cellulose membranes such as a Size 6 membrane having a molecular weight cut-off of 12-14 kDa, available from Medicell International Ltd., and tangential flow filtration (TFF) membranes, such as a Pellicon XL PLCCC 10 (50 cm<sup>2</sup>) or PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. In a preferred embodiment of the invention, the deprotected polypeptide or deprotected glatiramer is subjected to dialysis in water, followed by dialysis in aqueous acetic acid solution.

### III. Process Using an Amine or Ammonia.

In one embodiment of the invention, the process comprises:

- (a)<sup>2</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
- (b)<sup>2</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>2</sup> and a solvent, to form a product, preferably, the acid is admixed with a solution or suspension comprising the protected polypeptide and solvent; and
- (c)<sup>2</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia, and mixtures thereof, with the product formed in Step (b)<sup>2</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

In one embodiment of the invention, the process comprises:

- (a)<sup>2'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein said protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof;
- (b)<sup>2'</sup> admixing an acid with the protected glatiramer formed in Step (a)<sup>2'</sup> and a solvent, to form a product;
- (c)<sup>2'</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, with the product formed in Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water, to form a deprotected glatiramer; and
- (d)<sup>2'</sup> treating the deprotected glatiramer formed in Step (c)<sup>2'</sup> with acetic acid to form glatiramer acetate.

In the first deprotecting step, Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> of the process of the invention, an acid is admixed with the protected polypeptide formed in Step (a)<sup>2</sup> or protected glatiramer formed in Step (a)<sup>2'</sup>, and a solvent to form a product. Preferably, the acid is added to a mixture, i.e., a solution or suspension, comprising the protected polypeptide or protected glatiramer and a solvent. In Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> the protecting groups on the glutamic

acid moiety, i.e.,  $\gamma$ -*p*-methoxybenzyl group and/or  $\gamma$ -benzyl group, are removed. While not wishing to be bound by any particular theory, the present inventors believe that the acid also cleaves amide bonds of the protected polypeptide or protected glatiramer forming heterogenous polypeptide or glatiramer fragments.

Suitable acids include, but are not limited to, acetic acid, hydrochloric acid, hydrogen bromide, hydrogen fluoride, methane sulfonic acid, trifluoromethane sulfonic acid, phosphoric acid, trifluoroacetic acid and sulfuric acid. A mixture of acids may also be used. Preferred acids are selected from trifluoroacetic acid, a mixture of acetic acid and hydrochloric acid, a mixture of acetic acid and hydrogen bromide and a mixture of acetic acid and sulfuric acid. The acid may be added in the form of an aqueous solution.

The acid is preferably present in an amount of from about 0.1 wt. % to about 100 wt. %, more preferably from about 1 wt. % to about 10 wt. %, based on the weight of the protected polypeptide or protected glatiramer. Most preferably, the acid is present in an amount of from about 2 wt. % to about 6 wt. %, based on the weight of the protected polypeptide or protected glatiramer.

The temperature of the reaction medium during addition of the acid is preferably from about 10°C to about 40°C, more preferably 15°C to about 30°C. The acid is preferably added over a period of time from about 1 hour to about 30 hours, with stirring. Most preferably, the acid is added to the protected polypeptide or protected glatiramer at a temperature of about 25°C for a period of from about 1 hour to about 8 hours, with stirring.

The solvent used in the first deprotection step, Step (b)<sup>2</sup> and Step (b)<sup>2'</sup>, is selected from polar protic solvents and polar aprotic solvents. Preferably, the solvent used in Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> is selected from acetic acid, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane. A mixture of solvents may also be used. Most preferably, the solvent used in Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> is tetrahydrofuran or acetic acid.

The amount of solvent used in Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> is preferably from about 1-fold (wt.) to about 1,000-fold (wt.), more preferably, from about 10-fold (wt.) to about 500-fold (wt.), based on the amount of protected polypeptide or protected glatiramer which is used in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup>.

In the second deprotecting step, Step (c)<sup>2</sup> and Step (c)<sup>2'</sup> of the process of the invention, a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is admixed with the product formed in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or deprotected glatiramer.

Preferably, the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is added to a mixture, i.e., a solution or suspension, comprising the product formed in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water. The ammonia is in the form of NH<sub>3</sub> (aqueous) or NH<sub>3</sub> (gas). Preferably an aqueous solution of ammonia is used having a pH of about 7 to about 14. The addition of a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, to the product formed in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup> preferably removes the protecting group, such as *N*<sup>ε</sup>-trifluoroacetyl group, of the lysine moiety. Preferably the deprotected polypeptide is a deprotected glatiramer in the form of a free base.

The substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is preferably present in an amount of from about 1-fold (wt.) to about 1,000-fold (wt), more preferably from about 10-fold (wt.) to about 500-fold (wt), based on the total weight of the product of Step (b)<sup>2</sup> or Step (b)<sup>2'</sup> which is used in Step (c)<sup>2</sup> or Step (c)<sup>2'</sup>. Most preferably, the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is present in an amount of from about 50-fold (wt.) to about 150-fold (wt).

The substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is preferably admixed with a solution or suspension comprising the product formed in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water, at a temperature of from about 10°C to about 60°C, more preferably 15°C to about 40°C, for a period of time preferably from about 1 minute to about 60 hours, to form a deprotected polypeptide or deprotected glatiramer. Suitable solvents include, but are not limited to, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane. A preferred solvent is tetrahydrofuran.

More preferably, the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is admixed with a solution or suspension comprising the product formed in Step (b)<sup>2</sup> or Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water, at a temperature of from about 20°C to about 30°C, for a period of from about 1 hour to about 30 hours, to form a deprotected polypeptide or deprotected glatiramer. Unreacted diisopropylamine or ammonia, or any solvent or water, is preferably removed by evaporation or vacuum distillation.

Undesired low molecular weight polypeptide or glatiramer fragments, i.e., less than about 2 kDa, and high molecular weight polypeptide or glatiramer fragments, i.e., greater than about 40 kDa, are preferably removed by such methods as dialysis or diafiltration. Preferred membranes include Visking partially permeable cellulose membranes, such as a Size 6 membrane having a molecular weight cut-off of 12-14 kDa, available from Medicell International Ltd., and TFF membranes, such as a Pellicon XL PLCCC 10 (50 cm<sup>2</sup>) or PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. In a preferred embodiment of the invention, the deprotected polypeptide or deprotected glatiramer is subjected to dialysis in water, followed by dialysis in aqueous acetic acid solution.

#### IV. Process Using an Alkali or Alkaline Earth Metal Hydroxide, Carbonate or Hydrogencarbonate.

In one embodiment of the invention, the process comprises:

- (a)<sup>3</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
- (b)<sup>3</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>3</sup> and a solvent, to form a product, preferably, the acid is admixed with a solution or suspension comprising the protected polypeptide and solvent; and
- (c)<sup>3</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, with the product formed in Step (b)<sup>3</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

In one embodiment of the invention, the process comprises:

- (a)<sup>3'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein said protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof;
- (b)<sup>3'</sup> admixing an acid with a mixture comprising the protected glatiramer formed in Step (a)<sup>3'</sup> and a solvent, to form a product;
- (c)<sup>3'</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, with a mixture comprising the product formed in Step (b)<sup>3'</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected glatiramer; and
- (d)<sup>3'</sup> treating the deprotected glatiramer formed in Step (c)<sup>3'</sup> with acetic acid to form glatiramer acetate.

In the first deprotecting step, Step (b)<sup>3'</sup> and Step (b)<sup>3'</sup> of the process of the invention, an acid is admixed with the protected polypeptide formed in Step (a)<sup>3'</sup> or protected glatiramer formed in Step (a)<sup>3'</sup>, and a solvent to form a product. Preferably, the acid is added to a mixture, i.e., a solution or suspension, comprising the protected polypeptide or protected glatiramer and a solvent. In Step (b)<sup>3'</sup> and Step (b)<sup>3'</sup> the protecting groups on the glutamic acid moiety, i.e.,  $\gamma$ -*p*-methoxybenzyl group and/or  $\gamma$ -benzyl group, are removed. While not wishing to be bound by any particular theory, the present inventors believe that the acid also cleaves amide bonds of the protected polypeptide or protected glatiramer forming heterogenous polypeptide fragments.

Suitable acids include, but are not limited to, acetic acid, hydrochloric acid, hydrogen bromide, hydrogen fluoride, methane sulfonic acid, trifluoromethane sulfonic acid, phosphoric acid, trifluoroacetic acid and sulfuric acid. A mixture of acids may also be used. Preferred acids are selected from trifluoroacetic acid, a mixture of acetic acid and hydrochloric acid, a mixture of acetic acid and hydrogen bromide and a mixture of acetic acid and sulfuric acid. The acid may be added in the form of an aqueous solution.

The acid is preferably present in an amount of from about 0.1 wt. % to about 100 wt. %, more preferably from about 1 wt. % to about 10 wt. %, based on the weight of the protected polypeptide or protected glatiramer. Most preferably, the acid is present in an



amount of from about 2 wt. % to about 6 wt. %, based on the weight of the protected polypeptide or protected glatiramer.

The temperature of the reaction medium during addition of the acid is preferably from about 10°C to about 40°C, more preferably 15°C to about 30°C. The acid is preferably added over a period of time from about 1 hour to about 30 hours, with stirring. Most preferably, the acid is added to the protected polypeptide or protected glatiramer at a temperature of about 25°C for a period of from about 1 hour to about 8 hours, with stirring.

The solvent used in the first deprotection step, Step (b)<sup>3</sup> and Step (b)<sup>3'</sup>, is selected from polar protic solvents and polar aprotic solvents. Preferably, the solvent used in Step (b)<sup>3</sup> and Step (b)<sup>3'</sup> is selected from acetic acid, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane. A mixture of solvents may also be used. Most preferably, the solvent used in Step (b)<sup>2</sup> and Step (b)<sup>2'</sup> is tetrahydrofuran or acetic acid.

The amount of solvent used in Step (b)<sup>3</sup> and Step (b)<sup>3'</sup> is preferably from about 1-fold (wt.) to about 1,000-fold (wt.), more preferably, from about 10-fold (wt.) to about 500-fold (wt.), based on the amount of protected polypeptide or protected glatiramer which is used in Step (b)<sup>3</sup> or Step (b)<sup>3'</sup>.

In the second deprotecting step, Step (c)<sup>3</sup> and Step (c)<sup>3'</sup>, of the process of the invention, a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is admixed with the product formed in Step (b)<sup>3</sup> or Step (b)<sup>3'</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or deprotected glatiramer.

Preferably, the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is added to a mixture, i.e., a solution or suspension, comprising the product formed in Step (b)<sup>3</sup> or Step (b)<sup>3'</sup>, and water or a mixture of a solvent and water. The addition of a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, to the product formed in Step (b)<sup>3</sup> or Step (b)<sup>3'</sup> preferably removes the *N*<sup>ε</sup>-trifluoroacetyl group of the lysine moiety. Preferably the deprotected polypeptide is a deprotected glatiramer in the form of a free base.

The substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate and a hydrogencarbonate, and includes calcium hydroxide, lithium hydroxide, magnesium hydroxide, potassium hydroxide, barium hydroxide, sodium hydroxide, calcium carbonate, lithium carbonate, magnesium carbonate, potassium carbonate, sodium carbonate, calcium hydrogencarbonate, lithium hydrogencarbonate, magnesium hydrogencarbonate, potassium hydrogencarbonate and sodium hydrogencarbonate. More preferably, the substance used in Step (c)<sup>3</sup> or Step (c)<sup>3i</sup> is selected from the group consisting of sodium hydroxide, lithium hydroxide and potassium hydroxide.

The substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is preferably present in an amount of from about 1-fold (wt.) to about 1,000-fold (wt), more preferably from about 10-fold (wt.) to about 500-fold (wt), based on the total weight of the product of Step (b)<sup>3</sup> or Step (b)<sup>3i</sup> which is used in Step (c)<sup>3</sup> or Step (c)<sup>3i</sup>. Most preferably, the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is present in an amount of from about 50-fold (wt.) to about 150-fold (wt).

The substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is preferably admixed with a solution or suspension comprising the product formed in Step (b)<sup>3</sup> or Step (b)<sup>3i</sup>, and water or a mixture of a solvent and water, at a temperature of from about 10°C to about 60°C, more preferably 15°C to about 40°C, for a period of time preferably from about 1 minute to about 60 hours, to form a deprotected polypeptide or deprotected glatiramer. Suitable solvents include, but are not limited to, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane. A preferred solvent is tetrahydrofuran.

More preferably, the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is admixed with a solution or suspension comprising the product formed in Step (b)<sup>3</sup> or Step (b)<sup>3i</sup>, and water or a mixture of a solvent and water, at a temperature of from about 20°C to about 30°C, for a period of from about 1 hour to about 30 hours, to form a deprotected polypeptide or deprotected glatiramer. Unreacted alkali or alkaline earth metal hydroxide, carbonate, hydrogencarbonate or any solvent or water, is preferably removed by evaporation or vacuum distillation.

The amount of solvent or a mixture of a solvent and water which is used in Step (c)<sup>3</sup> or Step (c)<sup>3'</sup> is preferably from about 1-fold (wt.) to about 1,000-fold (wt.), more preferably, from about 10-fold (wt.) to about 500-fold (wt.), based on the total weight of the product of Step (b)<sup>3</sup> or Step (b)<sup>3'</sup> which is used in Step (c)<sup>3</sup> or Step (c)<sup>3'</sup>.

After the second deprotection step, Step (c)<sup>3</sup> and Step (c)<sup>3'</sup>, any layers are separated preferably in separatory funnel, or the solvent is preferably removed by evaporation or vacuum distillation. The deprotected polypeptide or deprotected glatiramer is preferably obtained as a solution in water.

Undesired low molecular weight polypeptide or glatiramer fragments, i.e., less than about 2 kDa, and high molecular weight polypeptide or glatiramer fragments, i.e., greater than about 40 kDa, are preferably removed by such methods as dialysis or diafiltration. Preferred membranes include Visking partially permeable cellulose membranes, such as a Size 6 membrane having a molecular weight cut-off of 12-14 kDa, available from Medicell International Ltd., and TFF membranes, such as a Pellicon XL PLCCC 10 (50 cm<sup>2</sup>) or PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. In a preferred embodiment of the invention, the deprotected polypeptide or deprotected glatiramer is subjected to dialysis in water, followed by dialysis in aqueous acetic acid solution.

The weight average molecular weight of the polypeptide or a pharmaceutically acceptable salt thereof which is prepared in accordance with the processes of the invention is preferably from about 2 kDa to about 30 kDa, more preferably from about 4.7 kDa to about 11 kDa, and most preferably from about 7 kDa to about 10 kDa, as determined by gas permeation chromatography (GPC). Preferably, the polypeptide or pharmaceutically acceptable salt thereof is substantially free of polypeptide fragments having a molecular weight of greater than about 40 kDa. Preferably, the polypeptide or pharmaceutically acceptable salt thereof is substantially free of polypeptide fragments having a molecular weight of less than about 2 kDa. As used herein, "substantially free" means less than about 5% by weight, more preferably less than about 2.5% by weight of the polypeptide prepared according to the process of the invention.

The polypeptide or pharmaceutically acceptable salt thereof, preferably glatiramer acetate, prepared according to the process of the invention may be formulated by conventional methods known in the art.

In one embodiment of the invention, the polypeptide or pharmaceutically acceptable salt thereof, which is prepared by the process of the invention, is dissolved or suspended in an acceptable pharmaceutical liquid vehicle, such as water, and the solution or suspension is injected into the body.

In one embodiment of the invention, the glatiramer acetate salt is dissolved in a mixture containing water and mannitol, and the solution is injected into the body.

Typically, the polypeptide or a pharmaceutically acceptable salt thereof is administered daily to patients suffering from multiple sclerosis, e.g., at a dosage of 20 mg of glatiramer acetate.

The following non-limiting examples illustrate further aspects of the invention.

Examples 1-7 relate to an acid hydrolysis process for preparing glatiramer acetate.

#### Example 1

##### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-alanine (860 mg, 7.5 mmol), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (600 mg, 2.3 mmol), *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine (1410 mg, 5.2 mmol) and *N*-carboxyanhydride of L-tyrosine (300 mg, 1.4 mmol) are placed in a three-neck flask. Distilled anhydrous dioxane (57 mL) is added. Diethylamine (3.4  $\mu$ L) is added. The resulting mixture is stirred mechanically for 24 hours at a temperature of approximately 22-25°C. The mixture is slowly poured into 100 mL deionized water and filtered under vacuum. The solid is kept under vacuum for 12 hours.

#### Example 2

##### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-alanine (430 mg, 3.75 mmol), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (300 mg, 1.15 mmol), *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine (705 mg, 2.6 mmol) and *N*-carboxyanhydride of L-tyrosine (150 mg, 0.7 mmol) are placed in a three-neck flask. Distilled anhydrous dioxane (28.5 mL) is added. Diethylamine (1.7  $\mu$ L) is added. The resulting mixture is stirred mechanically for 24 hours at a temperature of approximately 22-25°C. The mixture is slowly poured into 100 mL deionized water and filtered under vacuum. The solid is kept under vacuum for 12 hours.

### Example 3

Deprotecting the Protected Polypeptide Prepared in Example 1 with HBr/Acetic Acid.

The protected polypeptide prepared in Example 1 (200 mg) is suspended in 7 mL of 33% hydrogen bromide in acetic acid. The starting material slowly dissolves forming a red brown solution. The mixture is stirred for 17 hours at a temperature of approximately 22°C. The solution of HBr/acetic acid is evaporated to dryness using a rotary evaporator under reduced pressure. To this residue, 100 mL of water is added to dissolve the solid. The solution is placed in a Visking partially permeable cellulose membrane which is in the form of a tube, Size 6, having a molecular weight cut-off of 12-14 kDa. Size 6 tube has a diameter of 27/32 inches, 21.5 mm and a width of 32-34 mm. The tube is available from Medicell International Ltd. The tube containing the solution is stirred in a beaker of water. Polypeptide fragments having a molecular weight less than about 2 kDa are removed by osmosis from the dialysis tube. The tube is removed from the water and stirred in a beaker containing 0.3% acetic acid in water. The resulting product is removed from the tube and lyophilized to obtain glatiramer acetate as a pure white solid.

### Example 4

Deprotecting the Protected Polypeptide Prepared in Example 1 with HCl/Acetic Acid.

The protected polypeptide prepared in Example 1 (200 mg) is suspended in 20 mL of a mixture prepared of 9.4 mL concentrated hydrochloric acid adjusted to 20 mL with glacial acetic acid. The starting material slowly dissolves forming a slightly turbid solution. The mixture is stirred for 17 hours at a temperature of approximately 22°C. The solution of HCl/acetic acid is evaporated to dryness using a rotary evaporator under reduced pressure. To this residue, 100 mL of water is added to dissolve the solid. The solution is placed in a Visking partially permeable cellulose membrane which is in the form of a tube, Size 6, having a molecular weight cut-off of 12-14 kDa. Size 6 tube has a diameter of 27/32 inches, 21.5 mm and a width of 32-34 mm. The tube is available from Medicell International Ltd. The tube containing the solution is stirred in a beaker of water. Polypeptide fragments having a molecular weight less than about 2 kDa are removed by osmosis from the dialysis tube. The tube is removed from the water and stirred in a beaker containing 0.3% acetic acid in water. The resulting product is removed from the tube and lyophilized to obtain glatiramer acetate as a pure white solid.

### Example 5

Deprotecting the Protected Polypeptide Prepared in Example 2 with HCl/Acetic Acid.

The protected polypeptide prepared in Example 2 (200 mg) is suspended in 20 mL of a mixture prepared of 9.4 mL concentrated hydrochloric acid adjusted to 20 mL with glacial acetic acid. The starting material slowly dissolves forming a slightly turbid solution. The mixture is stirred for 17 hours at a temperature of approximately 22°C. The solution of HCl/acetic acid is evaporated to dryness using a rotary evaporator under reduced pressure. To this residue, 100 mL of water is added to dissolve the solid. The solution is placed in a Visking partially permeable cellulose membrane which is in the form of a tube, Size 6, having a molecular weight cut-off of 12-14 kDa. Size 6 tube has a diameter of 27/32 inches, 21.5 mm and a width of 32-34 mm. The tube is available from Medicell International Ltd. The tube containing the solution is stirred in a beaker of water. Polypeptide fragments having a molecular weight less than about 2 kDa are removed by osmosis from the dialysis tube. The tube is removed from the water and stirred in a beaker containing 0.3% acetic acid in water. The resulting product is removed from the tube and lyophilized to obtain glatiramer acetate as a pure white solid.

### Example 6

Relative molecular weight determination by GPC using a UV Detector.

#### GPC-UV Detector Conditions:

Eluent:	Phosphate buffer 0.05 M, pH 7.4, 5.6 g Na <sub>2</sub> HPO <sub>4</sub> , 116 g NaCl/4L water
Column:	PSS Suprema, 10 µm, 100 Å, 8 x 300 mm.
Temperature:	23°C
Pump:	TSP AS 3000 autosampler
Inj. – Vol.:	50 µL
Concentration:	about 2.0 mg/mL
Detector:	TSP UV2000 at 276 nm
GPC-Software:	PSS WinGPC Vers. 7.2

**Samples:** COPAXONE® (Lot # 5308036) wherein the mannitol has been removed by dialysis

COPAXONE® (Lot # 8040341) wherein the mannitol has been removed by dialysis

Sample A is glatiramer acetate prepared in Example 4 according to the process of the invention

Sample B is glatiramer acetate prepared in Example 5 according to the process of the invention

**Sample Preparation:** The weighted samples are dispersed in the eluent and allowed to stand for full hydration at room temperature for about 12 hours. The sample solution is filtered through a 1.0 filter unit (Schleicher & Schuell)

The test results for GPC are summarized in Table I.

**TABLE I**

<b>Sample</b>	<b>M<sub>w</sub> [kD]</b>
COPAXONE® (Lot # 5308036)	7.923
COPAXONE® (Lot # 8040341)	9.524
Sample A	8.551
Sample B	7.689

The test results in Table I clearly show that the weight average molecular weight of glatiramer acetate prepared by the process of the invention, Samples A and B, is 8.551 kDa and 7.689 kDa, respectively, and the weight average molecular weight of COPAXONE® is 7.923 kDa and 9.524 kDa, depending on the lot number.

#### Example 7

Relative molecular weight determination by gel permeation chromatography ( GPC ) using a refractive index detector.

#### GPC-RI Detector Conditions:

**Eluent:** Phosphate buffer 0.05 M, pH 7.4, 5.6 g Na<sub>2</sub>HPO<sub>4</sub>, 116 g NaCl/4L water

**Column:** PSS Suprema, 10 µm, 100 Å, 8 x 300mm

**Temperature:** 23°C

**Pump:** TSP AS 3000 autosampler

**Inj. – Vol.:** 50 µL

Concentration: about 2.0 mg/mL

Detector: Shodex RI 71

GPC-Software: PSS WinGPC Vers. 7.2

Samples: COPAXONE® (Lot # 5308036) wherein the mannitol has been removed by dialysis

COPAXONE® (Lot # 8040341) wherein the mannitol has been removed by dialysis

Sample A is glatiramer acetate prepared in Example 4 according to the process of the invention

Sample B is glatiramer acetate prepared in Example 5 according to the process of the invention

Sample Preparation: The weighted samples are dispersed in the eluent and allowed to stand for full hydration at room temperature for about 12 hours. The sample solution is filtered through a 1.0 filter unit (Schleicher & Schuell)

The test results for GPC with RI detector are summarized in Table II.

**TABLE II**

<b>Sample</b>	<b>M<sub>w</sub> [kD]</b>
COPAXONE® (Lot # 5308036)	8.663
COPAXONE® (Lot # 8040341)	9.641
Sample A	9.581
Sample B	8.224

The test results in Table II clearly show that the weight average molecular weight of glatiramer acetate prepared by the process of the invention, Samples A and B, is 9.581 kDa and 8.224 kDa, respectively, and the weight average molecular weight of COPAXONE® is 8.663 kDa and 9.641 kDa, depending on the lot number.

Examples 8-17 relate to a phase transfer process for preparing glatiramer acetate.

The present inventors have established a scale of 1 (low) to 5 (high) to quantify the influence of the alkaline compound (sodium hydroxide) on glatiramer formation. The test results are summarized in Table III.



Table III.

NaOH Conc. in 15 mL THF + 75 mg Glatiramer Free Base	Phase Separation	Recovery of Desired Molecular Weight Product from Aqueous Phase SM = 75 mg	Low Molecular Weight Polypeptide Formation	Product with Desired Molecular Weight Cut-off
0.1 N / 5 mL	No	1	1	1
0.25 N / 5 mL	1	5	1	5
0.25 N / 10 mL	1	5	1	5
0.5 N / 5 mL	3	3	2	4
0.5 N / 10 mL	4	3	3	3
0.5 N / 10 mL + 10 mL water	5	3	2	4
1.0 N / 5 mL	5	3	5	1

The results in Table III clearly show that 0.1-1.0 N sodium hydroxide solutions may be used to prepare the polypeptide or a pharmaceutically acceptable salt thereof products of the invention. The results in Table III also show that the use of a sodium hydroxide concentration of at least 0.25 N is preferred to facilitate phase separation of the organic phase and aqueous phase.

#### Example 8

##### Polymerization Step (a).

*N*-Carboxyanhydride of tyrosine (30 mg, 0.010 mm), *N*-carboxyanhydride of alanine (62 mg, 0.054 mm), *N*-carboxyanhydride of  $\gamma$ -benzyl glutamate (42 mg, 0.016 mm) and *N*-carboxyanhydride of *E*-*N*-trifluoroacetyllysine (100 mg, 0.037 mm), are placed in a single-neck flask with a magnetic stirrer. To this mixture is added 20 mL of dry tetrahydrofuran. A clear solution is obtained. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 48 hours. The tetrahydrofuran is removed by evaporation. Water (150 mL) is added to the residue and the stirring is continued. A white solid is obtained. The solid is filtered and dried in a dessicator under vacuum.

The yield is determined to be 154 mg.

### Example 9

#### Deprotecting Step (b).

Protected glatiramer acetate, 75 mg, prepared in Example 8 is transferred to a single-neck flask provided with a magnetic stirrer. Tetrahydrofuran (15 mL) is added to the flask and stirred. A clear solution is obtained. Aqueous solution of sodium hydroxide (8 mL) 0.5 N, is added. The addition of the sodium hydroxide solution results in the mixture becoming hazy. The mixture is stirred for 1 hour at 24-26°C. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous 1 N HCl solution to pH = 6.0 with stirring. The crude glatiramer free base solution is filtered using a nylon filter (0.2 micron Nylon Acrodisk).

### Example 10

Protected glatiramer acetate, 75 mg, prepared in Example 8 is transferred to a single-neck flask provided with a magnetic stirrer. Tetrahydrofuran (15 mL) is added to the flask and stirred. A clear solution is obtained. Aqueous solution of sodium hydroxide (10 mL) 0.25 N is added. The addition of the sodium hydroxide solution results in the mixture becoming hazy. The mixture is stirred for 16 hours at 25-26°C. The reaction mixture is centrifuged for 15 minutes. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous HCl solution to pH = 7-7.5 with stirring. Stirring is continued for an additional 30 minutes and the pH is determined to be pH = 8.0. The crude glatiramer free base solution is filtered.

### Example 11

#### Polymerization Step.

*N*-Carboxyanhydride of tyrosine (30 mg, 0.010 mm), *N*-carboxyanhydride of alanine (62 mg, 0.054 mm), *N*-carboxyanhydride of  $\gamma$ -benzyl glutamate (42 mg, 0.016 mm) and *N*-carboxyanhydride of *E*-*N*-trifluoroacetyllysine (100 mg, 0.037 mm), are placed in a single-neck flask with a magnetic stirrer. To this mixture is added 20 mL of dry dioxane. A clear solution is obtained. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 48 hours. To this, water (150 mL) is added slowly with stirring. A white solid is obtained. The solid is filtered and dried in a dessicator under vacuum.

The yield is determined to be 170 mg.

### Example 12

#### Deprotecting Step.

Protected glatiramer acetate, 75 mg, prepared in Example 11 is transferred to a single-neck flask provided with a magnetic stirrer. Dioxane (15 mL) is added to the flask and stirred. A clear solution is obtained. Aqueous solution of sodium hydroxide (10 mL) 0.5 N, is added. The addition of the sodium hydroxide solution results in the mixture becoming hazy. The mixture is stirred for 16 hours at 25-26°C. The reaction mixture is centrifuged for 15 minutes. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous HCl solution to pH = 7-7.5 with stirring. Stirring is continued for an additional 30 minutes and the pH is determined to be pH = 8.0. The crude glatiramer free base solution is filtered.

### Example 13

#### Deprotecting Step Conducted at Lower Temperature.

Protected glatiramer acetate, 75 mg, prepared in Example 12 is transferred to a single-neck flask provided with a magnetic stirrer. Tetrahydrofuran (15 mL) is added to the flask and the temperature of the solution is reduced to 0°C. Aqueous solution of sodium hydroxide (10 mL) 0.5 N, is added to the solution while maintaining a temperature of 0°C. The addition of the sodium hydroxide results in the solution becoming hazy. The solution is stirred for 3 hours at 0°C. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous HCl solution to pH = 7-7.5 with stirring at 0°C. Stirring is continued for an additional 30 minutes and the pH is determined to be approximately pH = 8.0. The crude glatiramer free base solution is filtered.

---

### Example 14

#### Deprotecting Step with Acetate Buffer.

Protected glatiramer acetate, 75 mg, prepared in Example 12 is transferred to a single-neck flask provided with a magnetic stirrer. Tetrahydrofuran (15 mL) is added to the flask and the temperature of the solution is reduced to 0°C. Aqueous solution of sodium hydroxide (10 mL) 0.5 N, and acetic acid (2 mL) is added to the solution while maintaining a temperature of 0°C and a pH = 12. The addition of the sodium hydroxide and acetic acid results in the solution becoming hazy. The solution is stirred for 3 hours at 0°C. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous HCl solution to pH = 7-7.5 with stirring at 0°C. Stirring is continued for an

additional 30 minutes and the pH is determined to be approximately pH = 8.0. The crude glatiramer free base solution is filtered.

#### Example 15

Protected glatiramer acetate, 75 mg, prepared in Example 8 is transferred to a single-neck flask provided with a magnetic stirrer. Dioxane (15 mL) is added to the flask and stirred. A clear solution is obtained. Aqueous solution of sodium hydroxide (10 mL) 0.25 N is added. The addition of the sodium hydroxide solution results in the mixture becoming hazy. The mixture is stirred for 16 hours at 25-26°C. The reaction mixture is centrifuged for 15 minutes. The formation of two phases is observed. The bottom layer is separated and acidified using dilute aqueous HCl solution to pH = 7-7.5 with stirring. Stirring is continued for an additional 30 minutes and the pH is determined to be pH = 8.0. The crude glatiramer free base solution is filtered.

#### Example 16

Diafiltration (Tangential Flow Filtration).

The crude glatiramer free base solution prepared in Example 9 is diluted to 120 with water. The dilute solution is first filtered through a 10 K diafiltration membrane, Pellicon XL, PLCCC 10 (50 cm<sup>2</sup>), available from Millipore, and then, filtered through a 10 K diafiltration membrane, Pellicon XL, PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. The concentrated solution obtained is lyophilized. A white powder is obtained.

#### Example 17

— Chromatographic Method of Purification of Glatiramer Acetate.

The crude glatiramer free base solution prepared in Example 9 is subjected to chromatographic separation. A column for gel filtration, FRACTOGEL TSK HW55 (600 x 26 mm) is prepared in a Superformance 26 Merck cartridge according to the manufacturer's instructions. The column is equilibrated with 0.2M ammonium acetate buffer pH 5.0, 30 mL of glatiramer free base solution samples (20 mg/mL, in 0.2 M ammonium acetate pH 5.0) are loaded on the column and fractions are collected every 10 minutes. A fraction having an average molecular weight of 7-8 KDa is isolated.

Examples 18-25 relate to a process for preparing glatiramer acetate using an alkali or alkaline earth metal hydroxide, carbonate, or a hydrogen carbonate.

### Example 18

#### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-tyrosine (207.19 mg, 1.0 mM), *N*-carboxyanhydride of L-alanine (620 mg, 5.4 mM), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (430 mg, 1.6 mM) and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine (1.01 g, 3.73 mM), are placed in a single-neck flask (100 mL) with a magnetic stirrer. To this mixture is added 40 mL of tetrahydrofuran. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 24 hours at a temperature of approximately 25°C. The mixture is slowly poured into 100 mL water while stirring. A solid is precipitated. The solid is filtered after 2 hours of stirring and washed with water. The solid is resuspended in 100 mL water and filtered. The solid is kept under vacuum for approximately 12 hours.

### Example 19

#### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-tyrosine (207.19 mg, 1.0 mM), *N*-carboxyanhydride of L-alanine (620 mg, 5.4 mM), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (430 mg, 1.6 mM) and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine (1.01 g, 3.73 mM), are placed in a single-neck flask (100 mL) with a magnetic stirrer. To this mixture is added 40 mL of dioxane. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 48 hours at a temperature of approximately 25°C. The mixture is slowly poured into 100 mL water while stirring. A solid is precipitated. The solid is filtered and washed with water. The solid is resuspended in 100 mL water and filtered. The solid is kept under vacuum for approximately 12 hours.

### Example 20

#### Cleavage of $\gamma$ -Benzyl Moiety from the Polypeptide Prepared in Example 2.

The protected polypeptide prepared in Example 19, 100 mg, is suspended in tetrahydrofuran (20 mL) and cooled in an ice water bath. Concentrated sulfuric acid, 4 mL, is added. The resulting clear solution is stirred for 20 hours at a temperature of approximately 25°C. The solvent, tetrahydrofuran, is removed by evaporation at 25°C to form a viscous liquid. Water, 50 mL, is added to the viscous liquid with stirring. A white precipitate forms which is filtered under vacuum and dried over phosphorous pentoxide under vacuum at 25°C

for approximately 12 hours in the dark. A white solid is obtained. The solid is filtered and dried in a dessicator under vacuum. The yield is determined to be 75 mg.

#### Example 21

Cleavage of *N*<sup>ε</sup>-trifluoroacetyl Moiety from the Polypeptide Prepared in Example 20.

The protected polypeptide prepared in Example 20, 75 mg, is dispersed in 12 mL of tetrahydrofuran, 4 mL of 0.5 M aqueous sodium hydroxide is added with stirring. The mixture is stirred for 3 hours at ambient temperature (approximately 22°C). The lower aqueous layer is separated and acidified with acetic acid to pH = 6.0.

#### Example 22

Cleavage of  $\gamma$ -Benzyl Moiety from the Polypeptide Prepared in Example 19.

The protected polypeptide prepared in Example 19, 1 g, is suspended in 50 mL of a mixture prepared of 47 mL concentrated HCl adjusted to 100 mL with glacial acetic acid. The starting material slowly dissolves forming a slightly turbid solution. The mixture is stirred for 18 hours at a temperature of approximately 22°C. The solution is poured into 1,000 mL of stirred water. A white precipitate is formed. The suspension is stirred for another 3 hours and then filtered. The product is washed with water and dried under vacuum at 50°C for approximately 17 hours.

#### Example 23

Cleavage of *N*<sup>ε</sup>-trifluoroacetyl Moiety from the Polypeptide Prepared in Example 22.

The protected polypeptide prepared in Example 22, 300 mg, is dispersed in 45 mL of tetrahydrofurane, 25 mL of 0.5 M aqueous sodium hydroxide is added with stirring. The mixture is stirred for 3 hours at ambient temperature (approximately 22°C). A clear, two-phase liquid system is formed. The lower aqueous layer is separated and acidified with acetic acid to pH = 6.0. The clear, colorless solution is filled into dialysis bags and dialyzed at ambient temperature once against 0.3% aqueous acetic acid, and then against water until a pH of 5.5 is reached. This solution is filtered and lyophilized to yield glatiramer acetate as a white solid.

#### Example 24

##### Diafiltration (Tangential Flow Filtration).

The glatiramer acetate solution prepared in Example 21 is adjusted to 120 mL with water to provide a 0.5–0.6 mg/mL concentration of the glatiramer acetate. The dilute solution is first filtered through a 30 K diafiltration membrane, Pellicon XL, PLCCC 10 (50 cm<sup>2</sup>), available from Millipore, and then, filtered through a 3 K diafiltration membrane, Pellicon XL, PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. The concentrated solution obtained is lyophilized to provide glatiramer acetate in solid form.

#### Example 25

##### Chromatographic Method of Purification of Glatiramer Acetate.

The glatiramer as extract of pH = 6 prepared in Example 23 is concentrated in vacuo to dryness and subjected to chromatographic separation. A column for gel filtration, FRACTOGEL TSK HW55 (600 x 26 mm) is prepared in a Superformance 26 Merck cartridge according to the manufacturer's instructions. The column is equilibrated with 0.2 M ammonium acetate buffer pH 5.0, 30 mL of glatiramer free base solution samples (20 mg/mL, in 0.2 M ammonium acetate pH 5.0) are loaded on the column and fractions are collected. A fraction having an average molecular weight of 7–10 kDa is isolated.

Examples 26–33 relate to a process for preparing glatiramer acetate using an amine or ammonia.

#### Example 26

##### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-tyrosine (207.19 mg, 1.0 mM), *N*-carboxyanhydride of L-alanine (620 mg, 5.4 mM), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (430 mg, 1.6 mM) and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine (1.01 g, 3.73 mM), are placed in a single-neck flask (100 mL) with a magnetic stirrer. To this mixture is added 40 mL of tetrahydrofuran. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 24 hours at a temperature of approximately 25 °C. The mixture is slowly poured into 100 mL water while stirring. A solid is precipitated. The solid is filtered after 2 hours of stirring and washed with water. The solid is resuspended in 100 mL water and filtered. The solid is kept under vacuum for approximately 12 hours.

### Example 27

#### Preparation of a Protected Polypeptide.

*N*-Carboxyanhydride of L-tyrosine (207.19 mg, 1.0 mM), *N*-carboxyanhydride of L-alanine (620 mg, 5.4 mM), *N*-carboxyanhydride of  $\gamma$ -benzyl L-glutamate (430 mg, 1.6 mM) and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine (1.01 g, 3.73 mM), are placed in a single-neck flask (100 mL) with a magnetic stirrer. To this mixture is added 40 mL of dioxane. Diethylamine (10  $\mu$ L) is added. The resulting mixture is stirred for 48 hours at a temperature of approximately 25°C. The mixture is slowly poured into 100 mL water while stirring. A solid is precipitated. The solid is filtered and washed with water. The solid is resuspended in 100 mL water and filtered. The solid is kept under vacuum for approximately 12 hours.

### Example 28

#### Cleavage of $\gamma$ -Benzyl Moiety from the Polypeptide Prepared in Example 27.

The protected polypeptide prepared in Example 27, 100 mg, is suspended in tetrahydrofuran (20 mL) and cooled in an ice water bath. Concentrated sulfuric acid, 4 mL, is added. The resulting clear solution is stirred for 20 hours at a temperature of approximately 25 °C. The solvent, tetrahydrofuran, is removed by evaporation at 25°C to form a viscous liquid. Water, 50 mL, is added to the viscous liquid with stirring. A white precipitate forms which is filtered under vacuum and dried over phosphorous pentoxide under vacuum at 25°C for approximately 12 hours in the dark. A white solid is obtained. The solid is filtered and dried in a dessicator under vacuum. The yield is determined to be 75 mg.

### Example 29

#### Cleavage of $\gamma$ -Benzyl Moiety from the Polypeptide Prepared in Example 27.

The protected polypeptide prepared in Example 27, 1 g, is suspended in 50 mL of a mixture prepared of 47 mL concentrated HCl adjusted to 100 mL with glacial acetic acid. The starting material slowly dissolves forming a slightly turbid solution. The mixture is stirred for 18 hours at a temperature of approximately 22°C. The solution is poured into 1,000 mL of stirred water. A white precipitate is formed. The suspension is stirred for another 3 hours and then filtered. The product is washed with water and dried under vacuum at 50°C for approximately 17 hours.



### Example 30

Evaluation of Amines to be Used to Cleave the *N*<sup>ε</sup>-trifluoroacetyl Moiety from the Polypeptide Prepared in Example 28.

The polypeptide prepared in Example 28, 75 mg, is suspended in 15 mL of water. An amine, 7 mL, is added to the suspension to provide an amine concentration of 3 M. A list of amines is provided in Table IV. Since a deprotected polypeptide is soluble in water, the reaction is monitored by the clarity of the solution. The results for each of the amines are summarized in Table IV.

**TABLE IV**

Amine	Result
Morpholine	No clear solution after 48 hours.
<i>N</i> -methyl-piperazine	No clear solution after 48 hours.
Dicyclohexylamine	No clear solution after 48 hours.
Di- <i>sec</i> -butylamine	No clear solution after 48 hours.
Pyrrolidine	No clear solution after 48 hours.
Diisopropylamine	Clear solution after about 1 hour.
Dipropylamine	No clear solution after 48 hours.
Isopropylamine	Clear solution after about 1.5 hours.
Methylamine (aqueous)	No clear solution after 48 hours.

The results in Table IV clearly show that a free base form of the polypeptide prepared in Example 28 is formed only upon the addition of diisopropylamine or isopropylamine. The results in Table I also show that dipropylamine, morpholine, *N*-methyl-piperazine, dicyclohexylamine, di-*sec*-butylamine, pyrrolidine, and methylamine failed to produce a free base form of the polypeptide.

Thus, applicants unexpectedly determined that in the second deprotection step of the process of the invention, Step (b), diisopropylamine and isopropylamine were the only amines that successfully removed the *N*<sup>ε</sup>-trifluoroacetyl group of the lysine moiety.

### Example 31

Cleavage of the *N*<sup>ε</sup>-trifluoroacetyl Moiety from the Polypeptide Prepared in Example 28.

The polypeptide prepared in Example 28, 75 mg, is suspended in 15 mL of water. Diisopropylamine, 7 mL, is added to the suspension to provide an amine concentration of 3 M. A milky-white solution becomes clear in approximately 1 hour and the clear solution is stirred at 25°C for 20 hours. The reaction mixture is evaporated at approximately 25°C to form crude glatiramer free base in the form of a viscous liquid. Fifty percent (50%) acetic

acid (15 mL) is added to the mixture and stirred for 30 minutes to form a glatiramer acetate solution.

### Example 32

Diafiltration (Tangential Flow Filtration).

The glatiramer acetate solution prepared in Example 31 is diluted to 120 mL with water. The dilute solution is first filtered through a 30 K diafiltration membrane, Pellicon XL, PLCCC 10 (50 cm<sup>2</sup>), available from Millipore, and then, filtered through a 3 K diafiltration membrane, Pellicon XL, PLCCC 5 (50 cm<sup>2</sup>), available from Millipore. The concentrated solution obtained is lyophilized to provide glatiramer acetate in solid form.

### Example 33

Chromatographic Method of Purification of Glatiramer Acetate.

The glatiramer acetate solution prepared in Example 31 is concentrated *in vacuo* to dryness and subjected to chromatographic separation. A column for gel filtration, FRACTOGEL TSK HW55 (600 x 26 mm) is prepared in a Superformance 26 Merck cartridge according to the manufacturer's instructions. The column is equilibrated with 0.2 M ammonium acetate buffer pH 5.0, 30 mL of glatiramer free base solution samples (20 mg/mL, in 0.2 M ammonium acetate pH 5.0) are loaded on the column and fractions are collected. A fraction having an average molecular weight of 7-10 kDa is isolated.

While the invention has been described with particular reference to certain embodiments thereof, it will be understood that changes and modifications may be made by those of ordinary skill within the scope and spirit of the following claims:

WHAT IS CLAIMED IS:

1. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:
  - (i) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof; and
  - (ii) adding an acid to the protected polypeptide formed in Step (i) to form a polypeptide or a pharmaceutically acceptable salt thereof, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety.
2. A process for preparing glatiramer acetate comprising:
  - (a) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate, and mixtures thereof;
  - (b) adding an acid to the protected glatiramer formed in Step (a) to form a glatiramer, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety; and
  - (c) treating the glatiramer formed in Step (b) with acetic acid to form glatiramer acetate.
3. The process according to Claim 1, wherein the acid is selected from the group consisting of acetic acid, hydrochloric acid, hydrogen bromide, hydrogen fluoride, methane sulfonic acid, trifluoromethane sulfonic acid, phosphoric acid, trifluoroacetic acid, sulfuric acid and mixtures thereof.

4. The process according to Claim 3, wherein the acid is a mixture of acetic acid and hydrochloric acid.
5. The process according to Claim 3, wherein the acid is a mixture of acetic acid and hydrogen bromide.
6. The process according to Claim 3, wherein the acid is a mixture of acetic acid and sulfuric acid.
7. The process according to Claim 3, wherein the acid is trifluoroacetic acid.
8. The process according to Claim 2, wherein the glatiramer acetate has a weight average molecular weight of about 2 kDa to about 30 kDa.
9. The process according to Claim 8, wherein the glatiramer acetate has a weight average molecular weight of about 4.7 kDa to about 11 kDa.
10. The process according to Claim 9, wherein the glatiramer acetate has a weight average molecular weight of about 7 kDa to about 10 kDa.
11. The process according to Claim 1, wherein the polypeptide is substantially free of polypeptide fragments having a molecular weight of greater than about 40 kDa.
12. The process according to Claim 1, wherein the polypeptide is substantially free of polypeptide fragments having a molecular weight of less than about 2 kDa.
13. The process according to Claim 1, wherein Step (ii) is conducted at a temperature of from about 10°C to about 40°C.
14. The process according to Claim 13, wherein Step (ii), is conducted at a temperature of from about 15°C to about 30°C.
15. The process according to Claim 14, wherein Step (ii), is conducted at a temperature of about 22°C to about 25°C.
16. The process according to Claim 1, wherein the initiator is diethylamine.

17. The process according to Claim 1, wherein the polar aprotic solvent is selected from the group consisting of tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide, dichloromethane and mixtures thereof.
18. The process according to Claim 17, wherein the polar aprotic solvent is 1,4-dioxane.
19. The process according to Claim 17, wherein the polar aprotic solvent is tetrahydrofuran.
20. The process according to Claim 1, wherein the acid is present in an amount of from about 0.1 weight percent (wt. %) to about 100 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.
21. The process according to Claim 20, wherein the acid is present in an amount of from about 1 wt. % to about 10 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.
22. The process according to Claim 21, wherein the acid is present in an amount of from about 2 wt. % to about 6 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.
23. A polypeptide or a pharmaceutically acceptable salt thereof which is prepared by a process comprising:
- (i) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof; and
  - (ii) adding an acid to the protected polypeptide formed in Step (i) to form a polypeptide or a pharmaceutically acceptable salt thereof, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety.

24. Glatiramer acetate which is prepared by a process comprising:
- (a) polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of protected L-glutamate and *N*-carboxyanhydride of *N*-*t*-butoxycarbonyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate, and mixtures thereof;
  - (b) adding an acid to the protected glatiramer formed in Step (a) to form a glatiramer, wherein said acid cleaves the  $\gamma$ -*p*-methoxybenzyl group from the glutamate moiety and the *N*-*t*-butoxycarbonyl group from the lysine moiety; and
  - (c) treating the glatiramer formed in Step (b) with acetic acid to form glatiramer acetate.
25. The glatiramer acetate which is prepared according to the process of Claim 24, wherein the glatiramer having a weight average molecular weight of less than 2 kDa is removed prior to Step (c).
26. The glatiramer acetate according to Claim 25, wherein the glatiramer having a weight average molecular weight of less than 2 kDa is removed by a method selected from the group consisting of dialysis and diafiltration.
27. The glatiramer acetate according to Claim 26, wherein the dialysis is initially conducted in water followed by dialysis in an aqueous acetic acid solution.
28. The glatiramer acetate which is prepared according to the process of Claim 24, wherein the process additionally comprises a lyophilization step after treating the glatiramer with acetic acid.
29. A method for treating a patient having an inflammatory, non-autoimmune central nervous system disease or alleviating the symptoms of such a disease, comprising administering a polypeptide or a pharmaceutically acceptable salt thereof prepared according to the process of Claim 1 to the patient in an amount and for a duration of time effective to treat the inflammatory, non-autoimmune central nervous system disease.

30. A method for treating a patient having an inflammatory, non-autoimmune central nervous system disease or alleviating the symptoms of such a disease, comprising administering glatiramer acetate prepared according to the process of Claim 2 to the patient in an amount and for a duration of time effective to treat the inflammatory, non-autoimmune central nervous system disease.
31. The method according to Claim 30, wherein the non-autoimmune central nervous system disease is multiple sclerosis.
32. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises treating a protected polypeptide with an aqueous solution of an alkali or alkaline earth metal hydroxide to form a polypeptide or a pharmaceutically acceptable salt thereof.
33. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:
- (a)<sup>1</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide; and
  - (b)<sup>1</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected polypeptide formed in Step (a)<sup>1</sup> to form a polypeptide or a pharmaceutically acceptable salt thereof.
34. A process for preparing glatiramer acetate comprising:
- (a)<sup>1'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a  $\gamma$ -benzyl L-glutamate L-glutamate and *N*-carboxyanhydride of N<sup>ε</sup>-trifluoroacetyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer;
  - (b)<sup>1'</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected glatiramer formed in Step (a)<sup>1'</sup> to form a glatiramer; and
  - (c)<sup>1'</sup> treating the glatiramer with acetic acid to form glatiramer acetate.
35. The process according to Claim 33, wherein Step (b)<sup>1</sup> is conducted at a temperature of from about -78°C to about 40°C.

36. The process according to Claim 35, wherein Step (b)<sup>1</sup>, is conducted at a temperature of from about -25°C to about 30°C.
37. The process according to Claim 36, wherein Step (b)<sup>1</sup>, is conducted at a temperature of from about -10°C to about 10°C.
38. The process according to Claim 37, wherein Step (b)<sup>1</sup>, is conducted at a temperature of from about 0°C.
39. The process according to Claim 32 wherein the pH is from about 13 to about 14.
40. The process according to Claim 32 which additionally comprises a buffer.
41. The process according to Claim 40, wherein the buffer is an acetate buffer and the pH is from about 8 to about 12.
42. The process according to Claim 32, wherein the alkali or alkaline earth metal hydroxide is selected from the group consisting of calcium hydroxide, lithium hydroxide, magnesium hydroxide, potassium hydroxide, sodium hydroxide and mixtures thereof.
43. The process according to Claim 42, wherein the alkali or alkaline earth metal hydroxide is sodium hydroxide.
44. The process according to Claim 32, wherein the alkali or alkaline earth metal hydroxide is present in an amount of from about 0.1 wt. % to about 400 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.
45. The process according to Claim 44, wherein the alkali or alkaline earth metal hydroxide is present in an amount of from about 10 wt. % to about 300 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.
46. The process according to Claim 45, wherein the alkali or alkaline earth metal hydroxide is present in an amount of from about 140 wt. % to about 260 wt. %, based on the total weight of the polypeptide or pharmaceutically acceptable salt thereof.



47. A polypeptide or a pharmaceutically acceptable salt thereof which is prepared by a process comprising treating a protected polypeptide with an aqueous solution of an alkali or alkaline earth metal hydroxide to form the polypeptide or pharmaceutically acceptable salt thereof, wherein said polypeptide comprises L-tyrosine, L-alanine, L-glutamate and L-lysine.
48. A polypeptide or a pharmaceutically acceptable salt thereof which is prepared by a process comprising:
- (a)<sup>1</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide; and
  - (b)<sup>1</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected polypeptide formed in Step (a)<sup>1</sup> to form a polypeptide or a pharmaceutically acceptable salt thereof.
49. Glatiramer acetate which is prepared by a process comprising
- (a)<sup>1'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a  $\gamma$ -benzyl L-glutamate L-glutamate and *N*-carboxyanhydride of N<sup>ε</sup>-trifluoroacetyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer;
  - (b)<sup>1'</sup> adding an aqueous solution of an alkali or alkaline earth metal hydroxide to the protected glatiramer formed in Step (a)<sup>1'</sup> to form a glatiramer; and
  - (c)<sup>1'</sup> treating the glatiramer formed in Step (b)<sup>1'</sup> with acetic acid to form glatiramer acetate.
50. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:
- (a)<sup>2</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
  - (b)<sup>2</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>2</sup> and a solvent, to form a product; and

(c)<sup>2</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia, and mixtures thereof, with the product formed in Step (b)<sup>2</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

51. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:

(a)<sup>2</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;

(b)<sup>2</sup> admixing an acid with a solution or suspension comprising the protected polypeptide formed in Step (a)<sup>2</sup> and a solvent, to form a product; and

(c)<sup>2</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia, and mixtures thereof, with the product formed in Step (b)<sup>2</sup>, and water or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.

52. The process according to Claim 50, wherein the protected L-lysine is *N*<sup>ε</sup>-trifluoroacetyl L-lysine.

53. The process according to Claim 50, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof.

54. The process according to Claim 53, wherein the protected L-glutamate is  $\gamma$ -benzyl L-glutamate.

55. A process for preparing glatiramer acetate comprising:

(a)<sup>2</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein said protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof;

- (b)<sup>2'</sup> admixing an acid with the protected glatiramer formed in Step (a)<sup>2'</sup> and a solvent, to form a product;
- (c)<sup>2'</sup> admixing a substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia, and mixtures thereof, with the product formed in Step (b)<sup>2'</sup>, and water or a mixture of a solvent and water, to form a deprotected glatiramer; and
- (d)<sup>2'</sup> treating the deprotected glatiramer formed in Step (c)<sup>2'</sup> with acetic acid to form glatiramer acetate.

56. The process according to Claim 50, wherein the solvent used in Step (b)<sup>2</sup> is selected from the group consisting of polar protic solvents, polar aprotic solvents and mixtures thereof.

57. The process according to Claim 56, wherein the solvent is selected from the group consisting of acetic acid, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane.

58. The process according to Claim 57, wherein the solvent is tetrahydrofuran.

59. The process according to Claim 57, wherein the solvent is acetic acid.

60. The process according to Claim 50, wherein the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is present in an amount of from about 1-fold (wt.) to about 1,000-fold (wt), based on the total weight of the product of Step (b)<sup>2</sup> which is used in Step (c)<sup>2</sup>.

61. The process according to Claim 60, wherein the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is present in an amount of from about 10-fold (wt.) to about 500-fold (wt).

62. The process according to Claim 61, wherein the substance selected from the group consisting of diisopropylamine, isopropylamine, ammonia and mixtures thereof, is present in an amount of from about 50-fold (wt.) to about 150-fold (wt).

63. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:
- (a)<sup>3</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
  - (b)<sup>3</sup> admixing an acid with the protected polypeptide formed in Step (a)<sup>3</sup> and a solvent, to form a product; and
  - (c)<sup>3</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate, and mixtures thereof, with the product formed in Step (b)<sup>3</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.
64. A process for preparing a polypeptide comprising L-tyrosine, L-alanine, L-glutamate and L-lysine, or a pharmaceutically acceptable salt thereof, wherein said process comprises:
- (a)<sup>3</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of a protected L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected polypeptide;
  - (b)<sup>3</sup> admixing an acid with a mixture comprising the protected polypeptide formed in Step (a)<sup>3</sup> and a solvent, to form a product; and
  - (c)<sup>3</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate, and mixtures thereof, with the product formed in Step (b)<sup>3</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected polypeptide or a pharmaceutically acceptable salt thereof.
65. The process according to Claim 63, wherein the protected L-lysine is *N*<sup>ε</sup>-trifluoroacetyl L-lysine.
66. The process according to Claim 63, wherein the protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof.

67. A process for preparing glatiramer acetate comprising:
- (a)<sup>3'</sup> polymerizing a mixture of *N*-carboxyanhydride of L-tyrosine, *N*-carboxyanhydride of L-alanine, *N*-carboxyanhydride of a protected L-glutamate and *N*-carboxyanhydride of *N*<sup>ε</sup>-trifluoroacetyl L-lysine, in a polar aprotic solvent in the presence of an initiator, to form a protected glatiramer, wherein said protected L-glutamate is selected from the group consisting of  $\gamma$ -*p*-methoxybenzyl L-glutamate,  $\gamma$ -benzyl L-glutamate and mixtures thereof;
  - (b)<sup>3'</sup> admixing an acid with a mixture comprising the protected glatiramer formed in Step (a)<sup>3</sup> and a solvent, to form a product;
  - (c)<sup>3'</sup> admixing a substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate, and mixtures thereof, with a mixture comprising the product formed in Step (b)<sup>3'</sup>, and a solvent or a mixture of a solvent and water, to form a deprotected glatiramer; and
  - (d)<sup>3'</sup> treating the deprotected glatiramer formed in Step (c)<sup>3'</sup> with acetic acid to form glatiramer acetate.
68. The process according to Claim 63, wherein the solvent used in Step (b)<sup>3</sup> is selected from the group consisting of polar protic solvents, polar aprotic solvents and mixtures thereof.
69. The process according to Claim 68, wherein the solvent is selected from the group consisting of acetic acid, tetrahydrofuran, ethyl acetate, dimethyl furan, dimethylformamide, 1,4-dioxane, dimethoxyethane, 1,2-dichloroethylene, dimethylsulfoxide and dichloromethane.
70. The process according to Claim 69, wherein the solvent is tetrahydrofuran.
71. The process according to Claim 69, wherein the solvent is acetic acid.
72. The process according to Claim 63, wherein the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate, and mixtures thereof, is present in an amount of from about 0.1 wt. % to about 10 wt. %, based on the total weight of solvent or mixture of a solvent and water, which is used in Step (c)<sup>3</sup>.

73. The process according to Claim 72 wherein the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is present in an amount of from about 0.1 wt. % to about 5 wt. %, based on the total weight of solvent or mixture of a solvent and water, which is used in Step(c)<sup>3</sup>.

74. The process according to Claim 63, wherein the solvent or mixture of a solvent and water, which is used in Step (c)<sup>3</sup>, is present in an amount from about 1-fold (wt.) to about 1000-fold (wt.), based on the total weight of the product of Step (b)<sup>3</sup>, which is used in Step (c)<sup>3</sup>.

75. The process according to Claim 74, wherein the solvent or mixture of a solvent and water, which is used in Step (c)<sup>3</sup>, is present in an amount from about 10-fold (wt.) to about 500-fold (wt.), based on the total weight of the product of Step (b)<sup>3</sup>, which is used in Step (c)<sup>3</sup>.

76. The process according to Claim 63, wherein the substance selected from the group consisting of an alkali or alkaline earth metal hydroxide, a carbonate, a hydrogencarbonate and mixtures thereof, is selected from the group consisting of calcium hydroxide, lithium hydroxide, magnesium hydroxide, potassium hydroxide, barium hydroxide, sodium hydroxide, calcium carbonate, lithium carbonate, magnesium carbonate, potassium carbonate, sodium carbonate, calcium hydrogencarbonate, lithium hydrogencarbonate, magnesium hydrogencarbonate, potassium hydrogencarbonate and sodium hydrogencarbonate.

77. The process according to Claim 76, wherein the substance is selected from the group consisting of sodium hydroxide, lithium hydroxide and potassium hydroxide.

## INTERNATIONAL SEARCH REPORT

International application No

PCT/US2005/039003

A. CLASSIFICATION OF SUBJECT MATTER  
C08G69/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07K C08G

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2004/043995 A (BRANTFORD CHEMICALS) 27 May 2004 (2004-05-27) the whole document	1-77
A	FR 2 133 926 A (YEDA RESEARCH & DEVELOPMENT) 1 December 1972 (1972-12-01) the whole document	1-77
A	WO 95/31990 A (YEDA RESEARCH & DEVELOPMENT) 30 November 1995 (1995-11-30) the whole document	1-77

☐ Further documents are listed in the continuation of Box C.☒ See patent family annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

23 March 2006

Date of mailing of the international search report

30/03/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Masturzo, P

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2005/039003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004043995	A	27-05-2004	AU 2003283152 A1 EP 1565486 A2	03-06-2004 24-08-2005
FR 2133926	A	01-12-1972	CA 1006508 A1 CH 570167 A5 DE 2217817 A1 GB 1393435 A IL 36670 A JP 57010132 B US 3849550 A	08-03-1977 15-12-1975 16-11-1972 07-05-1975 10-09-1974 25-02-1982 19-11-1974
WO 9531990	A	30-11-1995	AT 212857 T AU 2602195 A BR 9507758 A CA 2191088 A1 CN 1152874 A CN 1626238 A CZ 9603410 A3 DE 10299030 I1 DE 69525340 D1 DE 69525340 T2 DK 762888 T3 EE 9600170 A EP 0762888 A1 ES 2172586 T3 FI 964600 A HK 1008657 A1 HU 75672 A2 IL 113812 A JP 10500955 T JP 3715600 B2 JP 2003105000 A LU 90987 A9 MD 1443 F1 NL 300096 I1 NO 964976 A NZ 287335 A PL 317331 A1 PT 762888 T SK 149396 A3 TJ 392 B US 5800808 A	15-02-2002 18-12-1995 23-09-1997 30-11-1995 25-06-1997 15-06-2005 11-06-1997 23-01-2003 21-03-2002 19-09-2002 27-05-2002 16-06-1997 19-03-1997 01-10-2002 05-12-1996 30-08-2002 28-05-1997 29-06-2000 27-01-1998 09-11-2005 09-04-2003 25-02-2003 30-04-2000 01-10-2002 23-01-1997 25-02-1999 01-04-1997 31-07-2002 06-08-1997 13-10-2004 01-09-1998



**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**